

Visual and Hedonic Perception of Food Stimuli in Children with Autism Spectrum Disorders and their Relationship to Food Neophobia

Perception

0(0) 1–17

© The Author(s) 2019

Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/0301006619828300

journals.sagepub.com/home/pec**Anne-Claude Luisier**

Research Center in Neurosciences of Lyon, Claude Bernard University Lyon 1, France; Institute of Special Education, University of Fribourg, Switzerland; Brocoli Factory, Sion, Switzerland

Geneviève Petitpierre

Institute of Special Education, University of Fribourg, Switzerland

Annick Clerc Béro

Brocoli Factory, Sion, Switzerland

Anne-Raphaëlle Richoz, Junpeng Lao and Roberto Caldara

Department of Psychology, University of Fribourg, Switzerland

Moustafa Bensafi 

Research Center in Neurosciences of Lyon, Claude Bernard University Lyon 1, France

Abstract

The present study examined whether children with autism spectrum disorder (ASD) and typically developing (TD) children differed in visual perception of food stimuli at both sensorimotor and affective levels. A potential link between visual perception and food neophobia was also investigated. To these aims, 11 children with ASD and 11 TD children were tested. Visual pictures of food were used, and food neophobia was assessed by the parents. Results revealed that children

Corresponding author:

Moustafa Bensafi, Research Center in Neurosciences of Lyon, Claude Bernard University Lyon 1, Lyon, France.
Email: moustafa.bensafi@cnrs.fr

with ASD explored visually longer food stimuli than TD children. Complementary analyses revealed that whereas TD children explored more multiple-item dishes (vs. simple-item dishes), children with ASD explored all the dishes in a similar way. In addition, children with ASD gave more negative appreciation in general. Moreover, hedonic rating was negatively correlated with food neophobia scores in children with ASD, but not in TD children. In sum, we show here that children with ASD have more difficulty than TD children in liking a food when presented visually. Our findings also suggest that a prominent factor that needs to be considered is time management during the food choice process. They also provide new ways of measuring and understanding food neophobia in children with ASD.

Keywords

autism, sight, food neophobia, hedonic evaluation, eye tracking

Date Received: 30 November 2017; accepted: 14 January 2019

Introduction

Sensory processing plays a significant role in the acceptance and recognition of foods in typically developing (TD) children (Mennella, 2014; Mennella, Reiter, & Daniels, 2016; Zeinstra, Koelen, Kok, & de Graaf, 2007). Cognitive and affective processing of food-related stimuli are of most importance in dietary development in children (Bullinger, 2013; Doyen, 2011) and are thought to underlie certain difficulties in relation to food, most notably in individuals with autism spectrum disorder (ASD) who exhibit atypical sensory behavior. According to the American Psychiatric Association's Diagnostic and Statistical Manual, Fifth Edition (DSM-5), ASD is a neurodevelopmental disorder characterized by definite impairments in communication and social skills, narrow and intense interests, and stereotyped behaviors. But recent studies (Mottron, Belleville, Rouleau, & Collignon, 2014) have pointed that the phenotype of autism involves heterogeneous adaptive traits (not only disabilities but also strengths, for instance, perceptual strengths as well as sometimes overfunctioning). At least one in every 100 people has some form of autism (Baird et al., 2006; Kim et al., 2011). Autism affects more often boys than girls (4.5:1; Baio, 2014). Core traits of autism first emerge in infancy (Walsh, Elsabbagh, Bolton, & Singh, 2011) and then evolve over the lifespan.

A study performed on 95 children with ASD (Nadon, Feldman, Dunn, & Gisel, 2011a) reported a significant association between sensory issues (as measured with the Short Sensory Profile [SSP] questionnaire) and food-related behavioral problems (as measured with the Food Profile Questionnaire; Nadon et al., 2011a). The prevalence of eating issues is reported in 80% to 90% of children with ASD, compared with 13% to 50% of TD children (Cermak, Curtin, & Bandini, 2010; Fodstad & Matson, 2008; Matson, Fodstad, & Dempsey, 2009; Nadon, Feldman, Dunn, & Gisel, 2011b; Nadon, Feldman, & Gisel, 2013).

According to the typology established by Johnson, Foldes, Demand, and Brooks (2015), dietary behavior particular to children with ASD comprises certain forms of (a) food selectivity, which may be related to the texture or the way in which the food is presented (food selectivity based on type, texture, and presentation), (b) food refusal, and (c) behavioral problems during meals (e.g., "does not stay seated" ; Nadon et al., 2011b). Among the sensory modalities involved in the appreciation of food, sight plays a predominant role. Wadhera and Capaldi-Phillips (2014) detailed the physiological, cognitive, and emotional

mechanisms underlying visual influence on food acceptance, notably in children. In the same line, Aldridge, Dovey, and Halford (2009) reported that the visual categories created by children affect their willingness to taste food.

Atypical sensory functioning in individuals with ASD has been well documented in the last decade with regard to the visual system (Simmons et al., 2009). Compared with TD individuals, individuals with ASD present special features in the visual processing of social (Papagiannopoulou, Chitty, Hermens, Hickie, & Lagopoulos, 2014) and nonsocial stimuli (Mottron, Dawson, Soulières, Hubert, & Burack, 2006). For example, visual exploration is improved and intensified for nonsocial stimuli considered particularly attractive by individuals with ASD (South et al., 2008), such as visual images of vehicles, electronic objects, road signs, or sports equipment (Sasson, Turner-Brown, Holtzclaw, Lam, & Bodfish, 2008). It should be noted that this effect was also reported in TD children, although to a lesser degree. Taken together, these findings suggest that, in experimental settings, the type of stimulus chosen and its salience have an impact on the visual exploration strategies used by individuals with ASD (Saitovitch et al., 2013). The first aim of the present study was to extend these findings to food visual stimuli. To this end, visual processing by children with ASD and TD children was explored using an implicit objective sensorimotor measure, namely eye movement (Aim 1).

Emotions associated with dietary stimuli can also explain part of the relationship between sensory difficulties and altered food behavior. The role of sensory processing and its hedonic dimension (pleasure) in the regulation of eating behavior is well described in TD children (Cooke, 2007; Pliner, 2008; van der Horst, 2012). At a very early stage of development, TD children experience the pleasures of eating: sensory pleasure, pleasure in satiety, and relational pleasure (Bullinger, 2004, 2006, 2013; Holley, 2006). Psychological (Sander, Grandjean, & Scherer, 2005; Scherer, Schorr, & Johnstone, 2001) and biological (Rolls, 2012, 2015) models of affects postulate a strong link between emotion and cognition in the interpretation of environmental objects, including food. Rolls' (2012) model more specifically explores the role of emotion in acceptance or reject and decision-making, postulating that exteroceptive sensory factors perceived by the eater interact with internal signals of satiety in the orbitofrontal cortex to produce a hedonic state, linked with the reward value of food which subsequently causes appetite and leads to the act of eating.

The influence of emotion on food-related behavior has been reported in TD children, for instance, food rejection behaviors driven by visual stimuli perceived or felt as aversive (Brown & Harris, 2012; Lafraire, Rioux, Giboreau, & Picard, 2016; Martins & Pliner, 2006). In children with ASD, understanding how hedonic features extracted from sensory signals are processed may help clarify specificities of this population in relation to food. In nonfood areas, research showed significant differences between TD and ASD children in tasks involving perception and recognition of emotions (Nuske, Vivanti, & Disanayake, 2013; Uljarevic & Hamilton, 2013) and emotional regulation (Samson, Hardan, Lee, Phillips, & Gross, 2015; Samson, Hardan, Podell, Phillips, & Gross, 2015). The second aim of the present study was therefore to extend this concept to food stimuli by comparing hedonic perception of visual food stimuli between children with ASD and TD children (Aim 2).

Finally, a major hindrance in extending the diversity and acceptance of new foods is food neophobia, defined as a reluctance to consume or tendency to reject foods considered new by the eater (Dovey, Staples, Gibson, & Halford, 2008; Lafraire et al., 2016; Loewen & Pliner, 1999). Food neophobia was found to be associated with sensory experience in general (Aldridge et al., 2009; Shim, Kim, & Mathai, 2011) and sight in particular (Wadhera & Capaldi-Phillips, 2014), and with sensory functioning (Cooke, 2007) and

anxiety (Galloway, Lee, & Birch, 2003). Interestingly, a recent study showed that the level of food neophobia in children with ASD but not in TD children was negatively correlated with hedonic categorization of olfactory food stimuli (Luisier et al., 2015). The third aim of the present study was therefore to examine whether this relationship between food neophobia and hedonic perception in children with ASD is also observed for visual food stimuli (Aim 3).

Material and Methods

Participants

Eleven children with ASD (age range: 5.1–15.2 years; no female), and 11 TD children (age range: 5.1–15.2 years; 4 females) were included in the study and matched according to age (± 6 months). There was no significant difference between groups in terms of age (TD: 127.5 ± 28.5 months; ASD: 125.3 ± 29 months; Mann-Whitney $Z = -0.131$, *ns*, Cohen's $d = -.026$, 95% CI $[-.92, .86]$). Children with ASD were recruited from a specialized educational institute in Sion (Switzerland). They were diagnosed with ASD or pervasive developmental disorder according to the DSM-IV and were eligible for inclusion in the Swiss ASD Observatory. ASD diagnoses were confirmed on the Autism Diagnostic Observation Schedule (ADOS) (ADOS-1 or ADOS-2; Lord, Rutter, DiLavore, Risi, & Gotham, 1999; Lord et al., 2012). Children with ASD all had normal or corrected-to-normal vision. TD children were recruited from schools in the area of Fribourg, Switzerland, and had normal school performance, without any known behavioral or psychological disorder and normal or corrected-to-normal vision.

In both groups, food neophobia was assessed by the parents on a standard 10-items questionnaire (the French version of the Adapted Food Neophobia Scale) with good internal consistency (Reverdy, Chesnel, Schlich, Köster, & Lange, 2008). For each item, parents indicated to what extent the corresponding statement was true, on a 7-point scale from *Very true for my child* to *Not at all true for my child*. The 10 items were as follows: (a) My child is very particular about the foods he will eat (reversed scoring); (b) My child likes foods from different countries; (c) My child does not trust new foods (reversed scoring); (d) My child likes to try unusual foods; (e) When my child has the choice between different flavors for a certain food (e.g., ice-cream or sweets), it likes to choose a flavor that it does not know; (f) My child will try a dish, even if it does not know what is in it; (g) The foods my child know are sufficient for him (reversed scoring); (h) My child is willing to eat anything that is offered; (i) My child is afraid to eat things it has never had before (reversed scoring); and (j) My child will not taste a food when he does not know what it is (reversed scoring). For Questions 2, 4, 5, 6, and 8, the highest score (7 points) was given to the response “Very true for my child” and the lowest (1 point) to “Not at all true for my child”; for Questions 1, 3, 7, 9, and 10, the scores were reversed. The food neophobia score was obtained by summing the scores for the 10 questions (range: 10–70); the higher the score, the higher the neophobia grade.

The sensory profile was assessed with the French version (translation and publication by ECPA: Editions du Centre de Psychologie Appliquée) of the SSP (Dunn, 2010), filled out by the parents. The SSP is a standardized questionnaire that includes seven sections: (a) tactile sensitivity (seven items), (b) taste or smell sensitivity (four items), (c) movement sensitivity (three items), (d) underresponsive or seeks sensation (seven items), (e) auditory filtering (six items), (f) low energy or weak (six items), and (g) visual or auditory sensitivity (five items).

The internal reliability for the total test and sections on Cronbach's alpha ranges from 0.70 to 0.90 (Dunn, 1999).

The study had institutional review board approval (*Commission Cantonale Valaisanne d'Ethique Médicale*: IRB n° CCVEM 022/14). Parental consent was obtained for all children.

Stimuli

Stimuli comprised digital pictures of foods ($n=29$) regularly served to the ASD children in their institution and culturally familiar in Switzerland. Each picture consisted of a garnished white plate, placed against a light gray background (real diameter: 21 cm; on-screen diameter: 14 cm). The area and position of the food on the plate were checked using a stencil. The on-screen stimuli were presented at a viewing distance of 60 cm (at a visual angle of 13.36°), using the Psychophysics Toolbox (PTB-3: Brainard, 1997; Kleiner, Brainard, & Pelli, 2007) in a MATLAB environment (R2010a; The MathWorks, Natick, MA, USA). The images showed either a single-item dish (with only carrots, beans, peas, or pasta) or multiple-item dish (with a combination of two or three foods).

Procedure

Participants sat in a dimly lit room, at 60 cm from a 24" Dell LCD computer screen. The experiment consisted of two sessions. In both sessions, pictures displaying different dishes were presented one at a time in random order on the computer screen. The presentation of the picture was preceded by a white fixation-cross presented in the center of the screen for 2000 ms (Figure 1). Session 1 served to familiarize the children with the experimental setting and consisted in passively viewing the stimuli. Each picture was presented for 5000 ms. Session 2 (main experimental phase) involved a two-alternative forced-choice task in which the child had to decide whether they liked or disliked the food shown. In this task, the picture was presented until the participant answered. During this hedonic task, participants gave their answer using a Swiss computer keyboard in which the "S" key was labeled with a positive smiley and the "K" key with a negative smiley. All participants took part in both sessions.

Eye-Tracking Recording

Participants' eye movements were recorded with a SR Research Desktop-Mount EyeLink 2K eye-tracker using the EyeLink Toolbox extensions (Cornelissen, Peters, & Palmer, 2002). Gaze location on the screen was recorded for the dominant eye, with a sampling rate of 1000 Hz. A chin-and-forehead rest ensured stable head positioning; all subjects agreed to use the chin-and-forehead rest except from one child with ASD, who sat on the lap of his teacher who stabilized his chin with her hands. Nine-point calibration was conducted before each block; additional calibration was performed when necessary to ensure optimal recording quality.

Statistical Analyses

To test eye-movement differences between groups (Aim 1), five parameters were extracted from eye-movement data: number of fixations, mean fixation duration (s), sum of fixation durations (s), total path length (pixels), and mean path length (pixels). We applied a linear mixed model (LMM) to analyze children's eye movements. The group (children with ASD




Figure 1. Schematic representation of the procedure (viewing with yes or no hedonic response).

coded as 1 vs. TD children coded as 0) was considered as the main explanatory variable. The children and stimuli were considered as random-effect variables. As example, the LMM for mean fixation duration data can thus be expressed as follows:

$$\text{Mean fixation duration} \sim \text{Group} + (1|\text{Stim}) + (1|\text{Children})$$

For number of fixations, generalized linear mixed model (GLMM) for Poisson distributed data was fitted.

To analyze children's ratings of the pictures (two response levels: like and dislike), a binomial logistic mixed model was applied with group as the main explanatory variable and the observers and stimuli considered as random-effect variables.

All models were fitted with GLMER (package lme4 version 1.1-12; using R 3.2.2 - R Core Team, 2015).

To assess the relationship between hedonic ratings and neophobia scores in both ASD children and TD children (Aim 3), the mean hedonic ratings for all dishes were first calculated for each child. Linear regression was then performed, with neophobia score as independent variable and mean hedonic rating as dependent variable.

Note that in additional analyses, the age variable was included in the model (for eye-movement parameters and hedonic ratings).

Results

Questionnaires: Food Neophobia and SSP Scores in Both Groups. The range of food neophobia score was large for both groups (TD: 20–57; ASD: 17–70). There were significant differences in SSP global score (Mann-Whitney $Z = -2.61$, $p = .008$, Cohen's $d = 1.75$, 95% CI [.57, 2.92]) and for certain items: tactile sensitivity ($Z = -2.75$, $p = .005$, $d = 1.31$, 95% CI [.30, 2.32]) and “under-responsive/seek sensation” ($Z = -2.97$, $p = .002$, $d = 1.52$, 95% CI [.45, 2.59]) reflecting that ASD children showed more atypical scores than TD children. There were no significant differences for the other SSP items: movement sensitivity ($Z = -.88$, ns , $d = .68$, 95% CI [-.26, 1.63]), taste or smell sensitivity ($Z = -1.682$, ns , $d = .62$, 95% CI [-0.37, 1.61]), auditory filtering ($Z = -1.92$, $p = .063$, $d = .92$, 95% CI [-.07, 1.91]), low energy or weak ($Z = -.87$, ns , $d = .57$, 95% CI [-.36, 1.51]), or visual or auditory sensitivity ($Z = -.84$, ns , $d = .41$, 95% CI [-.55, 1.36]).

Table 1. Means and Standard Errors of the Five Eye-Movement Parameters and Linear Mixed Model Results of Eye-Movement Variables for Group Effect.

	ASD	TD	Group effect estimate ^a	SE	χ^2 ^b	df	$p(>\chi^2)$
Number of fixations	4.43 ± 2.89	3.31 ± 2.08	.26	.17	2.24	1	ns
Mean fixation duration	0.39 ± 0.24	0.40 ± 0.22	-.01	.05	.09	1	ns
Sum of fixation durations	1.39 ± 0.72	1.09 ± 0.49	.31	.16	4.03	1	.045
Total path length	646.5 ± 610.9	364.5 ± 367.8	281.35	139.07	4.06	1	.044
Mean path length	161.4 ± 82.2	132.2 ± 79.0	30.39	15.60	3.75	1	.053

Note. ASD = autism spectrum disorder; TD = typically developing; SE = standard error; ns = not significant.

^aChildren with ASD versus TD children.

^bDeviance difference to compare model including group variable and model not including group variable.

Aim 1: Compared with TD children, children with ASD showed different eye-movement activity when perceiving visual food items. Table 1 shows means and standard errors for the five eye-movement parameters and illustrates the main effects of group for all five eye-movement parameters. Significant group effects were observed for two parameters: Compared with TD children, children with ASD showed longer sum of fixation durations ($p = .045$); they also show longer total path length ($p = .044$). The group effect was close to significance for a third parameter: Compared with TD children, children with ASD showed longer mean path length ($p = .053$). When age variable was included in the model, the group effect remained statistically significant for the sum of fixation durations ($p = .044$) and it became marginally significant for the total path length ($p = .051$). The group effect remained marginally significant for the mean path length ($p = .052$).

Aim 2: Compared with TD children, children with ASD rated food images as less pleasant. Statistical analysis revealed a significant effect of group for hedonic ratings (group effect estimate = -2.32 ; $SE = .001$; $\chi^2 = 7.47$, $p = .006$): TD children rated the food images as more pleasant than children with ASD (Figure 2(a)). Note that when age variable was included in the model, the effect of group remained significant ($p = .005$).

Aim 3: An association between hedonic ratings and food neophobia score was observed in children with ASD. When looking at hedonic judgment distribution within each group, it should be noted that ratings were much more variable in children with ASD ($.290 \pm .05$) than in TD children ($TD: .44 \pm .02$). We then examined whether the observed hedonic variability was related to variability in food neophobia. First, a model including a food neophobia score-by-group interaction was computed: Although the interaction was not significant (interaction estimate = $-.012$; $t = -1.86$, $p = .078$, 95% CI $[-.025, .001]$, and Cohen's $f^2 = .19$), this analysis suggested (on a descriptive level) that the slope of the regression line was steeper for ASD children than for controls. To check this, we analyzed the groups separately. Results revealed a significant negative relationship between food neophobia score and hedonic rating in children with ASD (adjusted $R^2 = .374$, $F(1, 9) = 6.97$, $p = .027$): Children with ASD who rated the food images as more unpleasant were also those with higher food neophobia scores (Figure 2(b)). No significant relationship was observed between hedonic ratings and food neophobia score in TD children (adjusted $R^2 = -.11$, $F(1, 9) = .01$, ns; Figure 2(c)).




Figure 2. Hedonic ratings in TD and children with ASD and neophobia score. (a) TD children perceived the food images as more pleasant than children with ASD (** corresponds to a $p < .01$; bars and variations correspond to means and standard errors). (b) A significant negative relationship was observed between hedonic ratings and neophobia score in children with ASD. (c) No significant relationship was observed between hedonic ratings and neophobia score in TD children.

ASD = autism spectrum disorder; TD = typically developing.

Complementary Analysis 1: Dish containing a single versus multiple items is processed differently in TD versus ASD children. Scientific studies conducted in the general population show that children who are “picky and fussy” in their eating are less willing to consume mixed foods or foods they find difficult to identify (Lafraire et al., 2016). Moreover, the overall presentation (e.g., visual appearance and arrangement), the number of foods and their colors as they appear in the plate also play a critical role in the child’s visual appreciation of food (Zampollo, Kniffin, Wansink, & Shimizu, 2012): For instance, green foods are rejected more often than orange foods (Mennella, 2014; Mennella et al., 2016; Zeinstra et al., 2007). Given that it is currently unclear whether these effects are also present in children with ASD, we further asked this issue in a first complementary analysis and tested the hypothesis that the nature and diversity of stimuli (i.e., food containing a single vs. multiple items that vary in color and in visual presentation) influence the visual processing of the food in children with ASD. To test this, group differences in eye-movement as a function of diversity was explored by applying a linear mixed model (LMM) including group (children with ASD coded as 1 vs. TD children coded as 0) and diversity (dishes with multiple items coded as 1 vs. dishes with single item coded as 0). The group-by-item-diversity interaction was also examined. The children were considered as random-effect variables. As example, the LMM for mean fixation duration data can thus be expressed as follows:

$$\text{Mean fixation duration} \sim \text{Group} + \text{Diversity} + \text{Group} \times \text{Diversity} + (1|\text{Children})$$

Note that, for fixation number, GLMM for Poisson distributed data was fitted.

All models were fitted with GLMER (package lme4 version 1.1-12; using R 3.2.2 - R Core Team, 2015).

Table 2 shows means and standard errors for the five parameters. Table 3 illustrates the main effects of diversity and group and diversity-by-group interaction for all five eye-movement parameters. Effects of diversity were observed for four parameters: fixations’ number ($p < .001$), sum of fixation durations ($p = .040$), total path length ($p < .001$), and mean path length ($p < .001$). These effects reflected larger eye movement activity for

Table 2. Means and Standard Errors of the Five Eye-Movement Parameters in TD and ASD Participants as a Function of Stimulus Diversity (Single vs. Multiple).

	TD		ASD	
	Single	Multiple	Single	Multiple
Number of fixations	2.77 ± 1.19	3.56 ± 1.47	4.43 ± 2.41	4.47 ± 2.0
Mean fixation duration	.43 ± .13	.39 ± .13	.42 ± .16	.37 ± .09
Sum of fixation durations	1.00 ± .16	1.13 ± .28	1.46 ± .48	1.37 ± .47
Total path length	179.1 ± 160.2	384.9 ± 230.2	531.2 ± 457.8	633.6 ± 426.5
Mean path length	95.1 ± 40.7	146.3 ± 32.8	137.1 ± 46.4	171.1 ± 39.3

Note. ASD = autism spectrum disorder; TD = typically developing.

Table 3. Complementary Statistical Analyses of Eye-Movement Variables Using Linear Mixed Model for All Variables Except for Number of Fixations (Generalized Linear Model for Number of Fixations).

	Estimate ^a	SE	t ^b	p(> t)	χ ^{2c}	df	p(>χ ²)
Number of fixations					15.48	3	.001
Intercept	.96	.13	7.30	<.001			
Group	.44	.18	2.39	.017			
Diversity	.25	.07	3.57	<.001			
Group × Diversity	-.24	.09	-2.66	.008			
Mean fixation duration					6.36	3	.096
Intercept	.43	.04	10.92	<.001			
Group	-.01	.06	-.12	ns			
Diversity	-.04	.03	-1.55	ns			
Group × Diversity	-.01	.04	-.31	ns			
Sum of fixation durations					10.08	3	.018
Intercept	1.00	.12	8.38	<.001			
Group	.46	.17	2.74	.011			
Diversity	.13	.06	2.06	.040			
Group × Diversity	-.21	.09	-2.41	.016			
Total path length					28.50	3	<.001
Intercept	190.77	103.43	1.84	.077			
Group	357.34	145.94	2.45	.022			
Diversity	230.57	51.79	4.45	<.001			
Group × Diversity	-116.49	72.45	-1.61	ns			
Mean path length					47.08	3	<.001
Intercept	94.29	12.64	7.44	<.001			
Group	43.62	17.84	2.45	.020			
Diversity	52.02	9.26	5.62	<.001			
Group × Diversity	-19.12	12.95	-1.48	ns			

Note. SE = standard error; ns = not significant.

^aGroup effect: children with ASD versus TD children; diversity: multiple items versus single item.

^bz value and p(>|z|) for number of fixations variable.

^cDeviance difference to compare model including all variables and model including only intercept.

multiple- versus single-item food pictures in terms of number of fixations, sum of fixation durations, total path length, and mean path length. Effects of group were observed for the same variables: Compared with TD children, children with ASD showed a greater number of fixations ($p = .017$), longer sum of fixation durations ($p = .011$), longer total path length




Figure 3. Linear contrasts between *multiple* and *single* food displays independently for autistic and control observers using iMap4.

ASD = autism spectrum disorder; TD = typically developing; ns = not significant.

($p = .022$), and longer mean path length ($p = .020$). Finally, significant group-by-item-diversity interactions were observed for two eye-movement parameters: number of fixations ($p = .008$) and sum of fixation duration ($p = .016$) reflecting for TD children (but not for children with ASD) greater number of fixations and greater fixation durations for food with multiple items compared with food with single items.

Complementary Analysis 2: The spatial distribution of the fixation pattern is also modulated by diversity of the food image in TD versus children with ASD. To further explore the spatial distribution of the fixation pattern, we performed in a second complementary analysis a spatial mapping analysis of the fixation duration using iMap4 (Caldara & Mielle, 2011; Lao, Mielle, Pernet, Sokhn, & Caldara, 2015). iMap4 is a data-driven analysis framework for statistical fixation mapping using linear mixed model (LMM) and nonparametric statistics based on resampling (Lao et al., 2015). The fixation duration vector of each single trial was projected into a two-dimensional space according to the x - and y -coordinates of the fixation using iMap4. We then smoothed the raw fixation duration map using a two-dimensional Gaussian Kernel function with a sigma around 1° of visual angle. The single-trial smoothed fixation maps were normalized (z score) to better model the spatial pattern. To explore the multivariate structure in the resulting three-dimensional matrix (trials \times x -size \times y -size), we applied a representational dissimilarity matrix analysis of the smoothed fixation map basic on Mahalanobis distance (using the `rdmfixmap.m` function in iMap4). The average multivariate distance between two fixation maps within the same group of observers was computed for each stimulus (Figure 3). Moreover, each pixel in the smoothed fixation map was then fitted in iMap4 as the response variable using the following formula:

$$\text{Fixation Intensity}_{(x,y)} \sim \text{Group} + (1|\text{Stimuli}) + (1|\text{Observer}), 1 \leq x \leq x\text{Size}, 1 \leq y \leq y\text{Size}$$

The linear mixed models were fitted using Maximal Likelihood with the default iMap4 settings. Linear contrast of the model coefficients was performed as hypothesis testing, with




Figure 4. Descriptive example of fixation maps for one ASD and one TD observer when viewing two different food images displaying either single (left) or multiple items (right). ASD = autism spectrum disorder; TD = typically developing.

a bootstrap spatial clustering procedure threshold on the cluster mass as multiple comparison corrections (Lao et al., 2015).

In accordance with the first complementary analysis, results revealed a significant difference in the two-dimensional fixation maps between multiple- and single-food items in the control group (TD children) only: The control observers spent more time and explored more food images with multiple items compared with food images containing a single item (local maximum within the significant cluster: $F(1, 631) = 19.90, p < .001$; local minimum: $F(1, 631) = 3.86; p = .050$ cluster corrected). Children with ASD spent equal time on both types of food images (multiple and single items), using a similar fixation pattern (*ns*; Figures 3 and 4 for a descriptive example).

Discussion

The aims of our study were (a) to compare exploratory (e.g., sensorimotor) behavior between ASD and TD children in response to visual food stimuli, (b) to compare ASD and TD children in a task of assigning valence to these visual food stimuli, and (c) to study the relationship between this valence judgment and a behavioral attitude toward food (food neophobia).

Concerning the first two aims, results showed that, compared with TD children, children with ASD explored foods longer (in particular reflected by a significant higher oculomotor activity in terms of sum of fixation durations and total path length) before making a hedonic decision. Deeper exploration of a stimulus to judge its (un)pleasantness could be interpreted either as a difficulty experienced by the children with ASD in identifying their own emotional state, or as a difficulty in categorizing hedonically the stimulus. In one of the most recent empirical studies on alexithymia in children with ASD (25 children with ASD aged 8–13 years vs. 32 TD children aged 8–12 years), Griffin, Lombardo, and Auyeung (2015)

demonstrated a higher rate of alexithymia in children with ASD compared with the TD control group. In the field of olfaction, Legiša, Messinger, Kermol, and Marlier (2013), comparing how emotional responses to odors were reflected in peripheral nervous system responses (facial and autonomic responses) in children with ASD and matched controls (aged 8–14 years), reported that children with ASD were less likely to verbally express an affective state corresponding to their facial expression. The difficulty of evaluating the valence of stimuli could thus contribute to lengthening exploration duration. From a pedagogical point of view, this implies that children with ASD may need more time or more sensory explorations to decide whether or not they like food.

Regarding the relationship between hedonic categorization and attitude toward new food (third aim), we found a correlation between valence attribution and the degree of neophobia, specifically in children with ASD and not in TD children. The food neophobia score is an indicator of the risk of a child rejecting food that he or she considers to be new. The neophobia as measured at a given time reflects the relationship to foods based on previous dietary experiences (Dovey et al., 2008) and the cognitive features that characterize the eater at the time of measurement (Dovey, Aldridge, & Dignan, 2012; Lafraire et al., 2016). The observed correlation supports the hypothesis of a link between visual processing and the ability to appreciate food in children with ASD. Interestingly, the relationship between visual processing, neophobia, and hedonic processing is in line with observations in the olfactory domain; Luisier et al. (2015) found that less contrasted odor hedonic categorization was negatively correlated with food neophobia scores in children with ASD: The less they discriminated hedonically (especially for pleasant odors), the more neophobic they were.

Another result of interest revealed by complementary analyses was that children with ASD visually processed all dishes in a similar way, regardless of diversity (single or multiple items). In contrast, a more discriminating pattern was observed in TD children who visually inspected complex food stimuli more intensively than simple stimuli. Interestingly, this finding was observed with both quantitative (numbers of fixation and sum of fixation duration) and qualitative (spatial pattern) analyses. In children with ASD, this lesser differentiation in visual processing according to stimulus diversity is in agreement with the hypothesis of Mottron's perceptual model (Mottron et al., 2006). This model states that, in children with ASD, local processing of visual information (*bottom-up*—real immediate sensitive experience) is preferred (processing by default) and that the top-down (concept-driven) regulation (Hadjikhani et al., 2004; Müller & Nussbeck, 2008; Ropar & Mitchell, 2002), optional in this population, is never used when the local processing is felt to be more effective. In the dishes used as stimuli in the present study, each type of food had intrinsic nuances and irregularities (color, size, area, etc.) that could attract the attention of children with ASD and favor local processing (details apprehension) instead of descending conceptual processing (category apprehension). Conversely, in TD children, conceptual categorization (e.g., carrot, beans, peas, considered as types of items regardless of the manner in which they were prepared or presented) precedes the sensory visual processing of details favored by children with ASD: Food becomes complex by accumulation of conceptual categories rather than of visual details. In children with ASD, local processing may lengthen processing duration for stimuli already encountered but presented in a slightly different form, without difference in processing according to stimulus diversity (single vs. multiple items).

Regarding ratings of liking per se, children were asked to judge whether or not they liked the food presented on the screen. Children with ASD significantly more often attributed a negative valence to the food images than TD children. In TD children, the mechanisms underlying food rejection involve sensory and perceptual processing, categorization, and emotional evaluation (Lafraire et al., 2016). Recognition of food generally is a prerequisite

for acceptance. If not recognized, the food is usually rejected (Cooke, 2007; Dovey et al., 2008; Lafraire et al., 2016). An assumption that may be raised here is that similar mechanisms linking recognition to hedonic appreciation may also affect children with ASD.

Memories and in particular emotions evoked by the sight of food may also partly explain lack of appreciation. However, vegetables, which were particularly numerous in the present study, are generally less liked by children than other foods, especially because of their flavor and texture (Williams, Gibbons, & Schreck, 2005). Difficulty in texture acceptance often involves strong tactile oral defensiveness (=overresponsiveness, first described by Ayres [1964], which have been shown to correlate with food-related difficulties in many TD children [Smith, Roux, Naidoo, & Venter, 2005]). Interestingly, in the present study, the SSP showed a significant difference in tactile score between the two groups.

The present study was essentially an exploratory study with a relatively small sample. Although our study revealed preliminary—but new—findings, certain potential limitations need to be discussed. First, to maximize control of the nutritional context, as well as educational methods around meals and the known food repertoire of the children with ASD participating in the study, we chose to work with a single institution where children had at least one meal per working day. The total number of children participating in the study and the overrepresentation of boys depended on the number of children diagnosed with ASD in the institution and the parental approval. According to the sample size which was in fact low and the gender balance, it will therefore be necessary to replicate the study with a larger number of children. Second, the hedonic judgment requested in the study was only based on a binary choice between *I like* and *I don't like*. Despite the fact that this kind of instrument is recommended when researchers made the decision to collect information directly from individuals with special needs (Cuskelly, Moni, Lloyd, & Jobling, 2013; Hartley & Maclean, 2006) and despite the fact that all participants were able to make judgments using a binary choice effectively, we cannot discard the possibility that binary choices force the respondent to choose one or the other response and hide hesitations or ratings that may fall between these two options. Third, in order to avoid verbal means of collecting children's hedonic responses, we used a push-button device, which seemed to be suitable for most children. However, the children's understanding of the instructions was not checked; training in the use of the device might reduce any risk of misunderstanding. This device could be interesting in nonverbal children, to enable them to express their appreciation. Fourth, another major limitation deals with the specificity of the stimuli used for testing the influence of diversity on eye-movement parameters and categorization. The question whether our findings are specific to food stimuli or can be extended to other semantic domains (such as manufactured objects, like cars, toys . . .) should then be explored in future studies. Moreover, the images used for the study represented only foods belonging to children's daily lives; it would be interesting to study the visual exploration of foods unknown to children, in order to better characterize exploratory behavior in unfamiliar situations.

Conclusions

In conclusion, our study suggests that children with ASD have more difficulties than TD children in liking a food when presented visually. Such effect may be related to the diversity of the food, which might be more difficult to recognize due to intrinsic variation or varied presentation. Moreover, our study suggests that a prominent factor that needs to be considered is time management during the food choice process: Giving the child enough time to explore the food and then deciding on appreciation and acceptance can enable him or her to manage the sensory experience autonomously and could contribute to establishing food

familiarity. Finally, given the different processes (attention to details and categorization mechanisms) that seem to be involved in the hedonic processing of visual food stimuli, it would be interesting for future research to investigate whether there are salient sensory cues that could facilitate the recognition of a food by a child with ASD. Given the link between sensory processing and food neophobia, increasing valence for a sensory dimension that would make sense for children with ASD (e.g., smell and taste) could make the food carrying this dimension more familiar and favor its acceptance.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This research was supported by the Interdisciplinary mission of the CNRS (AUTON call, EMOTON project 2016–2018).

ORCID iD

Moustafa Bensafi  <http://orcid.org/0000-0002-2991-3036>

References

- Aldridge, V., Dovey, T. M., & Halford, J. C. G. (2009). The role of familiarity in dietary development. *Developmental Review, 29*, 32–44. doi:10.1016/j.dr.2008.11.001
- Ayres, A. J. (1964). Tactile functions. Their relation to hyperactive and perceptual motor behavior. *The American Journal of Occupational Therapy, 18*, 6–11.
- Baio, J. (2014). Prevalence of autism spectrum disorder among children aged 8 years—Autism and developmental disabilities monitoring network, 11 sites, United States, 2010. *Morbidity and Mortality Weekly Report. Surveillance Summaries, 63*, 1–21. doi:24670961
- Baird, G., Simonoff, E., Pickles, A., Chandler, S., Loucas, T., Meldrum, D., & Charman, T. (2006). Prevalence of disorders of the autism spectrum in a population cohort of children in South Thames: The Special Needs and Autism Project (SNAP). *Lancet, 368*, 210–215. doi:10.1016/S0140-6736(06)69041-7
- Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision, 10*, 433–436. doi:10.1163/156856897X00357
- Brown, S. D., & Harris, G. (2012). Rejection of known and previously accepted foods during early childhood: An extension of the neophobic response? *International Journal of Child Health and Nutrition, 1*, 72–81.
- Bullinger, A. (2004). *Le développement sensori-moteur de l'enfant et ses avatars—Un parcours de recherche* [The sensory-motor development of a child and its avatars - a research path]. Ramonville Saint-Agne: Éres.
- Bullinger, A. (2006). Approche sensorimotrice des troubles envahissants du développement [A sensorimotor approach to pervasive developmental disorders]. *Contraste, 25*, 1254–7689. doi:10.3917/cont.025.0125
- Bullinger, A. (2013). Les représentations de l'organisme dans l'autisme, perspective développementale [Representations of the organism in autism, a developmental perspective]. *La revue lacanienne, 14*, 173. doi:10.3917/lrl.131.0173
- Caldara, R., & Mielle, S. (2011). iMap: A novel method for statistical fixation mapping of eye movement data. *Behavior Research Methods, 43*, 864–878. doi:10.3758/s13428-011-0092-x

- Cermak, S. A., Curtin, C., & Bandini, L. G. (2010). Food selectivity and sensory sensitivity in children with autism spectrum disorders. *Journal of the American Dietetic Association, 110*, 238–246. doi:10.1016/j.jada.2009.10.032
- Cooke, L. J. (2007). The importance of exposure for healthy eating in childhood: A review. *Journal of Human Nutrition and Dietetics: The Official Journal of the British Dietetic Association, 20*, 294–301. doi:10.1111/j.1365-277X.2007.00804.x
- Cornelissen, F. W., Peters, E. M., & Palmer, J. (2002). The Eyelink Toolbox: Eye tracking with MATLAB and the Psychophysics Toolbox. *Behavior Research Methods, Instruments, & Computers, 34*, 613–617. doi:10.3758/BF03195489
- Cuskelly, M., Moni, K., Lloyd, J., & Jobling, A. (2013). Reliability of a method for establishing the capacity of individuals with an intellectual disability to respond to Likert scales. *Journal of Intellectual & Developmental Disability, 38*, 318–324. doi:10.3109/13668250.2013.832734
- Dovey, T. M., Aldridge, V. K., & Dignan, W. (2012). Developmental differences in sensory decision making involved in deciding to try a novel fruit. *British Journal of Health Psychology, 17*, 258–272.
- Dovey, T. M., Staples, P. A., Gibson, L. E., & Halford, J. C. G. (2008). Food neophobia and “picky/fussy” eating in children: A review. *Appetite, 50*, 181–193. doi:10.1016/j.appet.2007.09.009
- Doyen, C. (2011). Phases du développement du comportement alimentaire [Phases in the development of eating behaviour]. In M. Mouren, C. Doyen, M.-F. Le Heuzet, & S. Cook-Darzens (Eds.), *Les troubles du comportement alimentaire de l'enfant: Du nourrisson au pré-adolescent—Manuel diagnostique et thérapeutique* [Eating disorders in children: From infants to pre-adolescents – A diagnostic and therapeutic manual]. (pp. 3–11). Paris, France: Elsevier Masson SAS.
- Dunn, W. (1999). *The sensory profile manual*. San Antonio, TX: The Psychological Corporation.
- Dunn, W. (2010). Profil sensoriel mesurer l'impact des troubles sensoriels de l'enfant sur sa vie quotidienne (traduction française ecpa) [French translation of Dunn, W. (1999). The Sensory Profile Manual. San Antonio, Tex: The Psychological Corporation]. ECPA.
- Fodstad, J. C., & Matson, J. L. (2008). A comparison of feeding and mealtime problems in adults with intellectual disabilities with and without autism. *Journal of Developmental and Physical Disabilities, 20*, 541–550. doi:10.1007/s10882-008-9116-6
- Galloway, A. T., Lee, Y., & Birch, L. L. (2003). Predictors and consequences of food neophobia and pickiness in young girls. *Journal of the American Dietetic Association, 103*, 692–698. doi:10.1053/jada.2003.50134
- Griffin, C., Lombardo, M. V., & Auyeung, B. (2015). Alexithymia in children with and without autism spectrum disorders. *Autism Research, 9*, 773–780. doi:10.1002/aur.1569
- Hadjikhani, N., Chabris, C. F., Joseph, R. M., Clark, J., McGrath, L., Aharon, I., . . . Harris, G. J. (2004). Early visual cortex organization in autism: An fMRI study. *Neuroreport, 15*, 267–270. doi:10.1097/01.wnr.0000107523.38715.fa
- Hartley, S. L., & Maclean, W. E. (2006). A review of the reliability and validity of Likert-type scales for people with intellectual disability. *Journal of Intellectual Disability Research, 50*, 813–827. doi:10.1111/j.1365-2788.2006.00844.x
- Holley, A. (2006, December). Le plaisir, la mémoire et l'action [Pleasure, memory and action]. In *Actes du colloque de l'Institut Français pour la Nutrition, Plaisir, ami ou ennemi de notre alimentation?* [Pleasure, friend or enemy to the way we eat?] Institut Français pour la Nutrition.
- Johnson, C. R., Foldes, E., Demand, A., & Brooks, M. M. (2015). Behavioral parent training to address feeding problems in children with autism spectrum disorder: A pilot trial. *Journal of Developmental and Physical Disabilities, 27*, 591–607. doi:10.1007/s10882-015-9437-1
- Kim, Y. S., Leventhal, B. L., Koh, Y.-J., Fombonne, E., Laska, E., Lim, E.-C., Grinker, R. R. (2011). Prevalence of Autism spectrum disorders in a total population sample. *American Journal of Psychiatry, 168*, 904–912. doi:10.1176/appi.ajp.2011.10101532
- Kleiner, M., Brainard, D., & Pelli, D. (2007). What's new in Psychtoolbox-3? *Perception, 36*, (supplement), 14. 1–16.
- Lafraire, J., Rioux, C., Giboreau, A., & Picard, D. (2016). Food rejections in children: Cognitive and social/environmental factors involved in food neophobia and picky/fussy eating behavior. *Appetite, 96*, 347–357. doi:10.1016/j.appet.2015.09.008

- Lao, J., Mielle, S., Pernet, C., Sokhn, N., & Caldara, R. (2015). iMap 4: An Open Source Toolbox for the statistical fixation mapping of eye movement data with linear mixed modeling. *Journal of Vision, 15*, 1–17. doi:10.1167/15.12.793
- Legiša, J., Messinger, D. S., Kermol, E., & Marlier, L. (2013). Emotional responses to odors in children with high-functioning autism: Autonomic arousal, facial behavior and self-report. *Journal of Autism and Developmental Disorders, 43*, 869–879. doi:10.1007/s10803-012-1629-2
- Loewen, R., & Pliner, P. (1999). Effects of prior exposure to palatable and unpalatable novel foods on children's willingness to taste other novel foods. *Appetite, 32*, 351–366. doi:10.1006/appe.1998.0216
- Lord, C., Rutter, M., DiLavore, P., Risi, S., & Gotham, K. (1999). *Autism diagnostic observation schedule (ADOS) manual*. Los Angeles, CA: Western Psychological Services.
- Lord, C., Rutter, M., DiLavore, P. C., Risi, S., Gotham, K., & Bishop, S. L. (2012). *Autism diagnostic observation schedule, second edition (ADOS-2) manual (Part I): Modules 1–4*. Torrance, CA: Western Psychological Services.
- Luisier, A.-C., Petitpierre, G., Ferdenzi, C., Bérod, A. C., Giboreau, A., Rouby, C., & Bensafi, M. (2015). Odor perception in children with autism spectrum disorder and its relationship to food neophobia. *Frontiers in Psychology, 6*, 1–10. doi:10.3389/fpsyg.2015.01830
- Martins, Y., & Pliner, P. (2006). “Ugh! That’s disgusting!”: Identification of the characteristics of foods underlying rejections based on disgust. *Appetite, 46*, 75–85. doi:10.1016/j.appet.2005.09.001
- Matson, J. L., Fodstad, J. C., & Dempsey, T. (2009). The relationship of children's feeding problems to core symptoms of autism and PDD-NOS. *Research in Autism Spectrum Disorders, 3*, 759–766. doi:10.1016/j.rasd.2009.02.005
- Mennella, J. A. (2014). Ontogeny of taste preferences: Basic biology and implications for health. *The American Journal of Clinical Nutrition, 99*, 704S–711S. doi:10.3945/ajcn.113.067694
- Mennella, J. A., Reiter, A. R., & Daniels, L. M. (2016). Vegetable and fruit acceptance during infancy: Impact of ontogeny, genetics, and early experiences. *Advances in Nutrition, 7*, 2115–2195.
- Mottron, L., Belleville, S., Rouleau, G. A., & Collignon, O. (2014). Linking neocortical, cognitive, and genetic variability in autism with alterations of brain plasticity: The Trigger-Threshold-Target model. *Neuroscience and Biobehavioral Reviews, 47*, 735–752. doi:10.1016/j.neubiorev.2014.07.012
- Mottron, L., Dawson, M., Soulières, I., Hubert, B., & Burack, J. (2006). Enhanced perceptual functioning in autism: An update, and eight principles of autistic perception. *Journal of Autism and Developmental Disorders, 36*, 27–43. doi:10.1007/s10803-005-0040-7
- Müller, C. M., & Nussbeck, S. (2008). Do children with autism spectrum disorders prefer to match pictures based on their physical details or their meaning? *Journal of Mental Health Research in Intellectual Disabilities, 1*, 140–155. doi:10.1080/19315860801988244
- Nadon, G., Feldman, D. E., Dunn, W., & Gisel, E. (2011a). Association of sensory processing and eating problems in children with autism spectrum disorders. *Autism Research and Treatment, 2011*, 1–8. doi:10.1155/2011/541926
- Nadon, G., Feldman, D. E., Dunn, W., & Gisel, E. (2011b). Mealtime problems in children with autism spectrum disorder and their typically developing siblings: A comparison study. *Autism: The International Journal of Research and Practice, 15*, 98–113. doi:10.1177/1362361309348943
- Nadon, G., Feldman, D., & Gisel, E. (2013). Feeding issues associated with the autism spectrum disorders. In M. Fitzgerald (Ed.), *Recent advances in autism spectrum disorders—Volume I* (pp. 597–630). Rijeka, Croatia: InTech.
- Nuske, H. J., Vivanti, G., & Disanayake, C. (2013). Are emotion impairments unique to, universal, or specific in autism spectrum disorder? A comprehensive review. *Cognition and Emotion, 27*, 1042–1061.
- Papagiannopoulou, E. A., Chitty, K. M., Hermens, D. F., Hickie, I. B., & Lagopoulos, J. (2014). A systematic review and meta-analysis of eye-tracking studies in children with autism spectrum disorders. *Social Neuroscience, 9*, 610–632. doi:10.1080/17470919.2014.934966
- Pliner, P. (2008). Cognitive schemas: How can we use them to improve children's acceptance of diverse and unfamiliar foods? *British Journal of Nutrition, 99*, S2–S7.
- Reverdy, C., Chesnel, F., Schlich, P., Köster, E. P., & Lange, C. (2008). Effect of sensory education on willingness to taste novel food in children. *Appetite, 51*, 156–165. doi:10.1016/j.appet.2008.01.010

- Rolls, E. T. (2012). Taste, olfactory and food texture reward processing in the brain and the control of appetite. *The Proceedings of the Nutrition Society*, *71*, 488–501. doi:10.1017/S0029665112000821
- Rolls, E. T. (2015). Neural integration of taste, smell, oral texture, and visual modalities. In R. L. Doty (Ed.), *Handbook of olfaction and gustation* (pp. 1027–1047). Hoboken, NJ: John Wiley & Sons, Inc.
- Ropar, D., & Mitchell, P. (2002). Shape constancy in autism: The role of prior knowledge and perspective cues. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, *43*, 647–653. doi:10.1111/1469-7610.00053
- Saitovitch, A., Bargiacchi, A., Chabane, N., Phillipe, A., Brunelle, F., Boddaert, N., . . . Zilbovicius, M. (2013). Studying gaze abnormalities in autism: Which type of stimulus to use? *Open Journal of Psychiatry*, *03*, 32–38. doi:10.4236/ojpsych.2013.32A006
- Samson, A. C., Hardan, A. Y., Lee, I. A., Phillips, J. M., & Gross, J. J. (2015). Maladaptive behavior in autism spectrum disorder: The role of emotion experience and emotion regulation. *Journal of Autism and Developmental Disorders*, *45*, 3424–3432. doi:10.1007/s10803-015-2388-7
- Samson, A. C., Hardan, A. Y., Podell, R. W., Phillips, J. M., & Gross, J. J. (2015). Emotion regulation in children and adolescents with autism spectrum disorder. *Autism Research*, *8*, 9–18. doi:10.1002/aur.1387
- Sander, D., Grandjean, D., & Scherer, K. R. (2005). A systems approach to appraisal mechanisms in emotion. *Neural Networks*, *18*, 317–352. doi:10.1016/j.neunet.2005.03.001
- Sasson, N. J., Turner-Brown, L. M., Holtzclaw, T. N., Lam, K. S. L., & Bodfish, J. W. (2008). Children with autism demonstrate circumscribed attention during passive viewing of complex social and nonsocial picture arrays. *Autism Research*, *1*, 31–42. doi:10.1002/aur.4
- Scherer, K., Schorr, A., & Johnstone, T. (Eds.). (2001). *Appraisal processes in emotion: Theory, methods, research*. Oxford, England: Oxford University.
- Shim, J. E., Kim, J., & Mathai, R. A. (2011). Associations of infant feeding practices and picky eating behaviors of preschool children. *Journal of the American Dietetic Association*, *111*, 1363–1368. doi:10.1016/j.jada.2011.06.410
- Simmons, D. R., Robertson, A. E., McKay, L. S., Toal, E., McAleer, P., & Pollick, F. E. (2009). Vision in autism spectrum disorders. *Vision Research*, *49*, 2705–2739. doi:10.1016/j.visres.2009.08.005
- Smith, A. M., Roux, S., Naidoo, N. T. (Raj), & Venter, D. J. L. (2005). Food choices of tactile defensive children. *Nutrition*, *21*, 14–19. doi:10.1016/j.nut.2004.09.004
- South, M., Ozonoff, S., Suchy, Y., Kesner, R. P., McMahon, W. M., & Lainhart, J. E. (2008). Intact emotion facilitation for nonsocial stimuli in autism: Is amygdala impairment in autism specific for social information? *Journal of the International Neuropsychological Society*, *14*, 42–54. doi:10.1017/S1355617708080107
- Uljarevic, M., & Hamilton, A. (2013). Recognition of emotions in autism: A formal meta-analysis. *Journal of Autism and Developmental Disorders*, *43*, 1517–1526. doi:10.1007/s10803-012-1695-5
- van der Horst, K. (2012). Overcoming picky eating. Eating enjoyment as a central aspect of children's eating behaviors. *Appetite*, *58*, 567–574. doi:10.1016/j.appet.2011.12.019
- Wadhwa, D., & Capaldi-Phillips, E. D. (2014). A review of visual cues associated with food on food acceptance and consumption. *Eating Behaviors*, *15*, 132–143. doi:10.1016/j.eatbeh.2013.11.003
- Walsh, P., Elsabbagh, M., Bolton, P., & Singh, I. (2011). In search of biomarkers for autism: Scientific, social and ethical challenges. *Nature Reviews Neuroscience*, *12*, 603–612. doi:10.1038/nrn3113
- Williams, K. E., Gibbons, B. G., & Schreck, K. A. (2005). Comparing selective eaters with and without developmental disabilities. *Journal of Developmental and Physical Disabilities*, *17*, 299–309. doi:10.1007/s10882-005-4387-7
- Zampollo, F., Kniffin, K. M., Wansink, B., & Shimizu, M. (2012). Food plating preferences of children: The importance of presentation on desire for diversity. *Acta Paediatrica*, *101*, 61–66. doi:10.1111/j.1651-2227.2011.02409.x
- Zeinstra, G. G., Koelen, M. A., Kok, F. J., & de Graaf, C. (2007). Cognitive development and children's perceptions of fruit and vegetables; a qualitative study. *The International Journal of Behavioral Nutrition and Physical Activity*, *4*, 30. doi:10.1186/1479-5868-4-30